ORIGINAL PAPER



Seasonal variability of anthropogenic indices of PAHs in sediment from the Kuala Selangor River, west coast Peninsular Malaysia

Najat Masood · Normala Halimoon · Ahmad Zaharin Aris · Mohamad Pauzi Zakaria · Vahab Vaezzadeh · Sami M. Magam · Shuhaimi Mustafa · Masni Mohd Ali · Mehrzad Keshavarzifard · Sadeq Abdullah Abdo Alkhadher · Chui Wei Bong · Murad Ali Alsalahi

Received: 14 November 2017/Accepted: 8 May 2018/Published online: 25 May 2018 © Springer Science+Business Media B.V., part of Springer Nature 2018

Abstract Rapid increase in industrialization and urbanization in the west coast of Peninsular Malaysia has led to the intense release of petroleum and products of petroleum into the environment. Surface sediment samples were collected from the Selangor River in the west coast of Peninsular Malaysia during four climatic seasons and analyzed for PAHs and biomarkers (hopanes). Sediments were soxhlet extracted and further purified and fractionated through first and second step column chromatography. A gas

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s10653-018-0122-z) contains supplementary material, which is available to authorized users.

N. Masood · N. Halimoon (⊠) · A. Z. Aris · S. M. Magam · S. A. A. Alkhadher Environmental Forensics Laboratory, Faculty of Environmental Studies, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia e-mail: mala_upm@upm.edu.my

M. P. Zakaria · V. Vaezzadeh · C. W. Bong Institute of Ocean and Earth Sciences (IOES), University of Malaya, 50603 Kuala Lumpur, Malaysia

S. Mustafa

Halal Products Research Institute, Universiti Putra Malaysia, 4300 Serdang, Selangor, Malaysia

M. M. Ali

School of Environmental and Natural Resource Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia chromatography–mass spectrometry (GC–MS) was used for analysis of PAHs and hopanes fractions. The average concentrations of total PAHs ranged from 219.7 to 672.3 ng g⁻¹ dw. The highest concentrations of PAHs were detected at 964.7 ng g⁻¹ dw in station S5 in the mouth of the Selangor River during the wet inter-monsoonal season. Both pyrogenic and petrogenic PAHs were detected in the sediments with a predominance of the former. The composition of hopanes was homogeneous showing that petroleum hydrocarbons share an identical source in the study area. Diagnostic ratios of hopanes indicated that some of the sediment samples carry the crankcase oil signature.

M. Keshavarzifard

Department of Earth Sciences, Shiraz University, Shiraz, Iran

C. W. Bong

Laboratory of Microbial Ecology, Institute of Biological Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia

M. A. Alsalahi

Department of Marine Chemistry and Pollution, Faculty of Marine Science and Environment, Hodeidah University, Hodeidah, Yemen