

Two New Moray Eels of Genera *Diaphenchelys* and *Gymnothorax* (Anguilliformes: Muraenidae) from Taiwan and the Philippines

Wen-Chien Huang^{1,2,*}, David G. Smith³, Kar-Hoe Loh⁴, and Te-Yu Liao⁵

¹Doctoral Degree Program in Marine Biotechnology, National Sun Yat-sen University, Kaohsiung, Taiwan. *Correspondence: E-mail: t6733@hotmail.com (Huang)

²Doctoral Degree Program in Marine Biotechnology, Academia Sinica, Taipei, Taiwan.

³Smithsonian Institution, Museum Support Center, Suitland, MD, U.S.A. E-mail: smithd@si.edu (Smith)

⁴Institute of Ocean and Earth Sciences, University of Malaya, Kuala Lumpur, Malaysia. E-mail: khloh@um.edu.my (Loh)

⁵Department of Oceanography, National Sun Yat-sen University, Kaohsiung, Taiwan. E-mail: swp0117@gmail.com (Liao)

(Received 6 October 2020 / Accepted 7 March 2021 / Published xx April 2021)

Communicated by Hin-Kiu Mok

Two new moray eels of the genera *Diaphenchelys* and *Gymnothorax* from Taiwan and the Philippines are described. *Diaphenchelys laimospila* sp. nov. is described based on two specimens that represent the third species and a new geographic record of the genus. It can be distinguished from the other two congeners by numbers of cephalic sensory pores, vertebral formula, morphometric values, and the color pattern. *Gymnothorax pseudokidako* sp. nov. is a muraenid with a dark brown body covered by pale snowflake-like blotches. It differs from the most similar species *Gymnothorax kidako* (Temminck and Schlegel) by having a relatively short tail (50.5–53.0% vs. 52.9–56.4% of TL), more dentary teeth (17–26 vs. 16–20), fewer total vertebrae (134–139 vs. 137–143), and the absence of white margin on anal fin (vs. prominent white margin). These two new